

Abnormal vitamin D₃ metabolism in patients with primary Sjögren's syndrome

K Müller, P Oxholm, O H Sørensen, M Thymann, M Høier-Madsen, K Bendtzen

Abstract

Recent studies have suggested that vitamin D₃ may have an immunoregulatory role *in vitro*. The vitamin D₃ metabolism in 35 patients with primary Sjögren's syndrome was investigated by measuring blood concentrations of 1 α ,25-dihydroxyvitamin D₃ (1 α ,25-(OH)₂D₃) and 25-hydroxyvitamin D₃ (25-OHD₃), as well as phenotypes and blood concentrations of Gc globulin, the main vitamin D₃ binding protein in the blood. 25-OHD₃ concentrations were diminished, but those of 1 α ,25-(OH)₂D₃ were normal. There was no significant difference between the distribution of Gc phenotypes in the patients with primary Sjögren's syndrome and normal controls. Likewise, blood concentrations of Gc globulin corresponded to normal values. Among patients with increased concentrations of IgM rheumatoid factor there was a significant negative correlation between the serum titres of IgM rheumatoid factor and 25-OHD₃ concentrations.

Cytoplasmic receptors for 1 α ,25-dihydroxyvitamin D₃ (1 α ,25-(OH)₂D₃) have been identified in normal human monocytes as well as in activated, but not resting, human B and T lymphocytes.^{1,2} Moreover, pulmonary alveolar macrophages from patients with sarcoidosis are capable of 1 α -hydroxylation, which converts 25-OHD₃ to the biologically most active form—1 α ,25-(OH)₂D₃.³ Furthermore, interferon gamma or lipopolysaccharide can induce normal human monocytes and macrophages to exhibit 1 α -hydroxylase activity.⁴

In vitro studies suggest that 1 α ,25-(OH)₂D₃ has an immunoregulatory role. The hormone induces differentiation of monocytes⁵ and it blocks the production of interleukin-2⁶ and immunoglobulin,⁷ possibly by inhibiting the lymphocyte activating interleukin-1.⁸

Primary Sjögren's syndrome is a chronic autoimmune disease characterised by lymphoid infiltration of exocrine glands, especially lachrymal and salivary glands.⁹ Polyclonal B cell activation with hypergammaglobulinaemia¹⁰ and the presence of a broad range of autoantibodies, including IgM and IgA rheumatoid factors, as well as antinuclear antibodies, are characteristic of the syndrome.

The aim of this study was to investigate whether disturbed vitamin D₃ metabolism accompanies primary Sjögren's syndrome, by examining blood concentrations of vitamin D₃ and phenotypes of Gc globulin, the main vitamin D₃ binding protein in the blood.

Methods

Thirty five consecutive female patients with primary Sjögren's syndrome were studied. Their ages ranged from 27 to 69 years, mean 51 years. All patients fulfilled the Copenhagen diagnostic criteria for primary Sjögren's syndrome¹¹—that is, keratoconjunctivitis sicca evaluated by the Schirmer-I test, break-up time, and van Bijsterveld score, and xerostomia determined by unstimulated sialometry, salivary gland scintigraphy, and lower lip biopsy. The patients were not treated with immunosuppressive drugs other than non-steroidal anti-inflammatory drugs (n=4). The most common extraglandular manifestations were as follows: arthralgia/synovitis (n=18, 51%), myalgia/myositis (n=7, 20%), Raynaud's phenomenon (n=8, 23%) and pulmonary signs (n=7, 20%). Only a limited number of the patients showed other extraglandular manifestations such as purpura, interstitial nephritis, cutaneous vasculitis, intermittent non-infectious fever, and lymphoproliferative disorders. Serum samples were stored at -20°C until assay.

Serum 25-OHD₃ was examined by a radioassay using rachitic rat kidney cytosol as a binding protein. The normal mean (SD) was 28.2 (12.4) ng/ml (taken from healthy volunteers of both sexes, aged 18 to 60 years, from whom blood samples were obtained throughout the year; the mean therefore represents a sex, age, and seasonal average).¹² Serum 1 α ,25-(OH)₂D₃ was measured by a modification of a method described previously.¹³ Calf thymus receptors were used as a source of binding protein. Normal mean (SD) was 33.1 (15.3) pg/ml (healthy volunteers of both sexes aged 18-86 years). In these assays vitamin D₂ is measured in addition to vitamin D₃.

Ionised calcium was measured, as described previously.¹⁴ The normal range (pH=7.4) is 1.15-1.35 mmol/l.

Gc typing was performed by isoelectric focusing, followed by specific immunofixation on cellulose acetate.¹⁵ Gc levels were determined by rocket immunoelectrophoresis using purified human Gc globulin (Sigma Chemical Company, St Louis, Missouri, USA) as standard and polyclonal rabbit anti-Gc-antibody (Dakopatts, Denmark) (normal range 0.4-0.7 mg/ml).¹⁶

The concentrations of IgA and IgM rheumatoid factors were determined by an enzyme linked immunosorbent assay (ELISA), as described previously.¹⁷ Concentrations of IgM rheumatoid factor were determined by relating the content in the samples to an internal laboratory standard with a known concentration

Laboratory of Medical Immunology,
Medical Department
TTA, Rigshospitalet
State University
Hospital, Copenhagen,
Denmark
K Müller
P Oxholm
K Bendtzen

Department of Medicine,
Sundby Hospital,
Copenhagen, Denmark
O H Sørensen

Institute of Forensic
Genetics, University of
Copenhagen
M Thymann

Laboratory of
Autoimmune Serology,
State Serum Institute,
Copenhagen
M Høier-Madsen

Correspondence to:
Dr Müller.

Accepted for publication
21 September 1989

of IgM rheumatoid factor expressed in international units (IU) (normal value <8 IU/ml). IgA rheumatoid factor concentrations were related to an internal laboratory standard serum and the content expressed in arbitrary units (AU) (normal value <25 AU/ml).

Titres of circulating antinuclear antibodies of the IgG class (determined by an indirect fluorescence technique) and concentrations of plasma immunoglobulins were recorded at the same time.

The Pearson correlation coefficient and the χ^2 of test was used. A level of 5% was regarded as significant.

Results

The blood concentrations of $1\alpha,25-(\text{OH})_2\text{D}_3$ in patients with Sjögren's syndrome (median 26.0, Q₁ 19.0, Q₃ 42.0 pg/ml) were normal (mean (SD) 33.1 (15.3) pg/ml). In contrast, the concentrations of 25-OHD₃ were reduced (median 9.0, Q₁ 5.0, Q₃ 16.0 ng/ml), (normal mean: 28.2 (12.4) ng/ml). Undetectable concentrations of 25-OHD₃ were found in nine patients, and six of these had serum $1\alpha,25-(\text{OH})_2\text{D}_3$ concentrations lower than 15 pg/ml (figure).

There was no significant difference between the distribution of Gc phenotypes in the patients with primary Sjögren's syndrome and in 1674 controls ($\chi^2=1.65$) (table). The serum levels of

Gc globulin (median 0.6 mg/ml, Q₁ 0.55, Q₃ 0.65) corresponded to normal values (0.4–0.7 mg/ml serum).

Increased concentrations of IgM rheumatoid factor were found in 25 (72%) patients, with a median concentration of 118 IU/ml (Q₁ 41, Q₃ 189). Increased concentrations of IgA rheumatoid factor were found in 21 (60%) of the patients, with a median concentration of 92 AU/ml (Q₁ 55, Q₃ 105). Among patients with increased concentrations of IgM rheumatoid factor there was a significant negative correlation ($r=-0.42$, $p<0.05$) between the titres of IgM rheumatoid factor and the concentration of 25-OHD₃ but not those of $1\alpha,25-(\text{OH})_2\text{D}_3$. This was, however, not found within the group of IgA rheumatoid factor positive patients.

Twenty four (69%) of the patients exhibited antinuclear antibodies. The concentration of $1\alpha,25-(\text{OH})_2\text{D}_3$ and 25-OHD₃ within this group did not differ significantly from the group of patients as a whole.


Sixteen (46%) had hypergammaglobulinaemia (IgG, IgA, IgM). The median IgG level was 14.9 g/l (Q₁ 12.7, Q₃ 25.2 g/l). Neither immunoglobulin levels nor the presence of extraglandular manifestations correlated with the vitamin D₃ concentrations.

Ionised calcium concentration in 12 patients from whom additional serum samples were available (median 1.19 mmol/l, Q₁ 1.14, Q₃ 1.23) was normal (1.15–1.35 mmol/l).

Gc phenotype distribution in 35 patients with primary Sjögren's syndrome

Number	Gc phenotypes					
	IF	IS	2	IF-IS	2-IF	2-IS
Observed	0	13	3	8	3	8
Expected*	0.9	11.4	2.5	6.4	3.0	10.8

*Calculated from Gc prevalence determined in 1674 unrelated Danish adults.



Distribution of $1\alpha,25$ -dihydroxyvitamin D₃ ($1\alpha,25-(\text{OH})_2\text{D}_3$) and 25-hydroxyvitamin D₃ (25-OHD₃) in 35 patients with primary Sjögren's syndrome. ($1\alpha,25-(\text{OH})_2\text{D}_3$: normal mean (SD) is 33.1 (15.3) pg/ml; 25-OHD₃: normal mean (SD) is 28.2 (12.4) ng/ml. ▲ = patients with undetectable concentrations of 25-OHD₃.

Discussion

Primary Sjögren's syndrome was chosen for two reasons to examine the hypothesis that vitamin D metabolism is changed in patients with systemic autoimmune diseases. Firstly, polyclonal B cell activation is a pronounced and convenient marker of disturbed immune regulation in this disease. Secondly, patients with primary Sjögren's syndrome are normally not treated with immunosuppressive drugs that may affect metabolic rate. To our knowledge, this study is the first investigation of vitamin D metabolism in patients with primary Sjögren's syndrome. We found that the serum concentrations of 25-OHD₃ were significantly below those of normal subjects in most patients with primary Sjögren's syndrome. In contrast, the median concentration of $1\alpha,25-(\text{OH})_2\text{D}_3$ was normal. Serum immunoglobulins and immune complexes do not interfere with the vitamin D₃ assay.

It is questionable whether the changed vitamin D metabolism affects immune reactivities and contributes to the polyclonal B cell activation that is characteristic of these patients. Because $1\alpha,25-(\text{OH})_2\text{D}_3$ is, immunologically, the most active vitamin D metabolite in vitro, it is conceivable that an abnormal concentration of this compound might result in changed immune function.¹⁻⁷ Low concentrations (≤ 18 pg/ml), however, were only found in eight of the 35 patients and did not correlate with the titres of immunoglobulin or autoantibodies. 25-OHD₃ has been found to be about a thousand times less active than $1\alpha,25-(\text{OH})_2\text{D}_3$ in vitro,^{6,8,18} and it has a higher affinity for Gc globulin.¹⁹ As

the normal blood concentrations of 25-OHD₃ are about a thousand times higher than those of 1 α ,25-(OH)₂D₃ the observed low concentrations of 25-OHD₃ might have affected immune reactivity in these patients. It is interesting, therefore, that a highly significant correlation between the concentration of circulating 25-OHD₃ and IgG titres has been reported.²⁰ This was not found in our study, but the negative correlation between 25-OHD₃ concentrations and IgM rheumatoid factor suggests that vitamin D₃ metabolism may be related to immunopathological processes in primary Sjögren's syndrome.

What might be the cause of the observed low 25-OHD₃ concentrations? One possibility might be an impaired ability of Gc globulin to bind D₃ vitamins. Thus certain phenotypes of Gc globulin have been associated with other immunoinflammatory diseases, such as rheumatoid arthritis,²¹ multiple sclerosis,²² and liver disease.²³ Eales and colleagues also recently reported an association between different Gc alleles and the susceptibility to, and the clinical manifestation of, human immunodeficiency virus (HIV) infection.²⁴ Subsequent studies, were, however, unable to confirm this finding.²⁵ In our patients with primary Sjögren's syndrome the distribution of Gc phenotypes did not differ significantly from that in the normal population. Moreover, the serum concentrations of Gc globulin were normal. Therefore, changed Gc globulin binding of vitamin D₃ is unlikely to account for the low 25-OHD₃ concentrations found in patients with primary Sjögren's syndrome.

Low concentrations of vitamin D₃ may also result from severely impaired liver function²⁶ but our patients had no signs of liver disease which might have explained the concentrations of 25-OHD₃ we found.

As the primary aim of this investigation was to study the vitamin D₃ metabolism, and because the concentrations of ionised calcium were normal, the parathyroid hormone and calcitonin metabolism were not likely to have been disturbed and were therefore not studied.

Finally, polyclonal B cell activation may itself have contributed to the low concentrations of 25-OHD₃, because 25-OHD₃ is metabolised in activated monocytes and consumed by activated lymphocytes through binding to vitamin D receptors of these cells.¹⁷

In conclusion, we found severely diminished blood concentrations of 25-OHD₃ in patients with primary Sjögren's syndrome. It is unclear whether the changed vitamin D₃ metabolism is related to the immunopathology of this disease or is merely an epiphenomenon.

- 1 Tsoukas C D, Deftos C J, Provedini D M, Manolagas S C. 1,25 Dihydroxyvitamin D₃ receptors in human leukocytes. *Science* 1983; 221: 1181-3.
- 2 Bhalla A K, Amento E F, Clemens T L, Holick M F, Krane S M. Specific high-affinity receptors for 1,25-dihydroxyvitamin D₃ in human peripheral blood mononuclear cells: presence in monocytes and induction in T lymphocytes following activation. *J Clin Invest* 1983; 57: 1308-10.
- 3 Adams J S, Sharma O P, Mercedes A, Singer F R. Metabolism of 25-hydroxyvitamin D₃ by cultured pulmonary alveolar macrophages in sarcoidosis. *J Clin Invest* 1983; 72: 1856-60.
- 4 Reichel H, Koeffler H P, Bishop J E, Norman A W. 25-Hydroxyvitamin D₃ metabolism by lipopolysaccharide-stimulated normal human macrophages. *J Clin Endocrinol Metab* 1987; 64: 1-9.
- 5 Abe E, Miyaura C, Sakagami H, et al. Differentiation of mouse myeloid leukemia cells induced by 1,25-dihydroxyvitamin D₃. *Proc Natl Acad Sci USA* 1981; 78: 4990-4.
- 6 Tsoukas C D, Provedini D M, Manolagas S C. 1,25-Dihydroxyvitamin D₃: a novel immunoregulatory hormone. *Science* 1984; 224: 1438-40.
- 7 Lemire J M, Adams J S, Sakai R, Jordan S C. 1,25-Dihydroxyvitamin D₃ suppresses proliferation and immunoglobulin production by normal human peripheral blood mononuclear cells. *J Clin Invest* 1984; 74: 657-61.
- 8 Müller K, Svenson M, Bendtzen K. 1,25-Dihydroxyvitamin D₃ and MC 903 are potent inhibitors of human interleukin-1 in vitro. *Immunol Lett* 1988; 17: 361-6.
- 9 Youinou P, Pennec P. Immunopathological features of primary Sjögren's syndrome. *Clin Exp Rheumatol* 1987; 5: 173-84.
- 10 Whaley K, Buchanan W W. Sjögren's syndrome and associated diseases. In: Parker C W, ed. *Clinical immunology*. Philadelphia: Saunders, 1980: 632-66.
- 11 Manthorpe R, Oxholm P, Prause J U, Schiødt M. The Copenhagen criteria for Sjögren's syndrome. *Scand J Rheumatol* 1986; (suppl 61): 19-21.
- 12 Lund B, Sørensen O H. Measurement of 25-hydroxyvitamin D in serum and its relation to sunshine, age and vitamin D intake in the Danish population. *Scand J Clin Lab Invest* 1979; 39: 23-30.
- 13 Lund B, Lund B, Sørensen O H. Measurement of circulating 1,25-dihydroxyvitamin D in man. Changes in serum concentrations during treatment with 1 α -hydroxycholecalciferol. *Acta Endocrinol (Copenh)* 1979; 91: 338-50.
- 14 Fogh-Andersen N, Christiansen T F, Komarmy L, Siggaard-Andersen O. Measurement of free calcium ion in capillary blood and serum. *Clin Chem* 1978; 24: 1545-52.
- 15 Thymann M. Gc subtypes determined by agarose isoelectrofocusing. Distribution in Denmark and application to paternity cases. *Hum Hered* 1981; 31: 214-21.
- 16 Putnam F W, ed. *Plasma proteins*. Vol 1. 2nd ed. New York: Academic Press, 1975; 60.
- 17 Vejtorp M, Høier-Madsen M, Halberg P. Enzyme-linked immunosorbent assay for determination of IgM rheumatoid factor. *Scand J Rheumatol* 1979; 8: 63-70.
- 18 Rigby W F C, Stacy T, Fanger M W. Inhibition of T lymphocyte mitogenesis by 1,25-dihydroxyvitamin D₃ (Calcitriol). *J Clin Invest* 1984; 74: 1451-5.
- 19 Franceschi R T, Simpson R O, DeLuca H F. Binding proteins for vitamin D metabolites: serum carriers and intracellular receptors. *Arch Biochem Biophys* 1981; 210: 1-13.
- 20 Sedrani S H. Correlation between concentrations of humoral antibodies and vitamin D nutritional status: a survey study. *Eur J Clin Nutr* 1988; 42: 243-8.
- 21 Papiha S S, Pal B. Gc (vitamin D binding protein) subtypes in rheumatoid arthritis. *Hum Genet* 1985; 70: 278-80.
- 22 Gonates N K, Greene M I, Waksman B H. Genetic and molecular aspects of demyelination. *Immunology Today* 1986; 7: 121-6.
- 23 Brown I R F, Carter N D, Sood A. Vitamin D binding globulin phenotypes in liver disease. *Clin Chim Acta* 1979; 95: 75-82.
- 24 Eales C J, Nye K E, Parkin J M, Weber J N, Forster S M, Harris J R W. Association of different allelic forms of group specific component with susceptibility to and clinical manifestation of human immunodeficiency virus infection. *Lancet* 1987; i: 999-1002.
- 25 Thymann M, Dickmeiss E, Sveigaard A, Pedersen C, Bygbjerg I, Faber V. AIDS and Gc protein. *Lancet* 1987; i: 1378.
- 26 Lund B, Sørensen O H, Hilden M, Lund B. The hepatic conversion of vitamin D in alcoholics with varying degrees of liver affection. *Acta Med Scand* 1977; 202: 221-4.